



Pharmasin®

200 MG/ML SOLUTION FOR INJECTION



Composition:

Pharmasin® 200 mg/ml solution contains 200.000 IU tylosin/ml.

Target species:

Cattle, sheep, goats, pigs.

Indications:

Cattle (adult)

Treatment of respiratory infections, metritis caused by Gram-positive micro-organisms, mastitis caused by *Streptococcus spp.*, *Staphylococcus spp.* or *Mycoplasma* and interdigital necrobacillosis, i.e. panaritum or foot rot.

Calves

Treatment of respiratory infections and necrobacillosis.

Pigs

Treatment of enzootic pneumonia, haemorrhagic enteritis, erysipelas and metritis.
Treatment of arthritis caused by *Mycoplasma* and *Staphylococcus spp.*

Sheep and goats

Treatment of respiratory infections, metritis caused by Gram-positive microorganisms, mastitis caused by Gram+ microorganisms or *Mycoplasma spp.****

Contra indications:

Do not administer to horses. Intramuscular injection can be fatal in chickens and turkeys.
Do not administer to animals with a known hypersensitivity to tylosin or other macrolides.
In cattle maximum injection volume per injection site should not exceed 15 ml.

Dosages and treatment periods:

Intramuscular or slow intravenous injection (only in cattle)

Cattle	Sheep and goat	Pigs
5-10 mg/tylosin/kg bodyweight per day during 3 days	10 mg tylosin/kg bodyweight per day during 3 days	5-10 mg tylosin/kg bodyweight per day during 3 days
2.5 to 5 ml solution for injection per 100 kg bodyweight	5 ml solution for injection per 100 kg bodyweight	2.5 to 5 ml solution for injection per 100 kg bodyweight

Withdrawal period(s):

Meat and offal

- Cattle: 28 days
- Sheep: 42 days
- Goats: 42 days
- Pigs: 14 days
- Milk: 108 hours

Shelf-life and storage:

- Shelf life of the veterinary medicinal product as packaged for sale: 24 months.
- Shelf life after first opening of the immediate packaging: 28 days.

Packaging:

The product is presented in a **50/ 100/ 250 ml** glass vials.*



* Check with your local Huvepharma representative which pack sizes are available in your region as this may vary.

**Pharmasin® 200 mg/ml solution for injection is following the authorized EU SPC (available at request).

*** Indications listed above are not necessarily authorized in all countries. Please consult the local label for exact indications and posology.