

## Research Article

# The Effect of the Targeted Application of Commercially Available Non-Antibiotic Additives in the Poultry Industry on Antibiotic Usage and Biological Performance under Commercial Field Conditions

Maro Simitopoulou<sup>1\*</sup>, Alain Bousquet-Mésidou<sup>2</sup>, Kyriacos Angastiniotis, George Kefalas<sup>3</sup>, Savvas Dimitriadis<sup>3</sup>, Sotiris Papisolomontos and Miltiades Ioannou

<sup>1</sup>VitaTrace nutrition Ltd, Propylaion 18, Strovolos Industriale state, 2033 Nicosia, Cyprus

<sup>2</sup>Intheres, Universitéde Toulouse, Inrae, Env, Toulouse, France

<sup>3</sup>Nuevo SA, Schimatari Votias, 32009, Greece

## Abstract

Modern poultry production systems often constrain natural poultry behavior and may result in health problems which in turn necessitate the use of antibiotics under veterinary prescription. Within the European Union (EU) there is increased pressure to limit if not prohibit the use of antibiotics for prophylactic as well as metaphylactic purposes in food producing animals. This situation has encouraged increased attention to be placed on farm biosecurity and stock management improvement as well as the appraisal, development and in-field use of non-antibiotic alternatives (NAA) such as organic acids, exogenous feed enzymes, prebiotics and probiotics, as well as a variety of phyto-genic products. The selection and usage of most NAAs commercially available as feed and/or water additives is often

\*Corresponding author: Simitopoulou M, VitaTrace Nutrition Ltd, Propylaion 18, Strovolos Industrial Estate, 2033 Nicosia, Cyprus, Email: maro.simitopoulou@gmail.com

**Citation:** Simitopoulou M, Bousquet-Mésidou A, Angastiniotis K, Kefalas G, Dimitriadis S, et al. (2024) The Effect of the Targeted Application of Commercially Available Non-Antibiotic Additives in the Poultry Industry on Antibiotic Usage and Biological Performance under Commercial Field Conditions 8: 059.

**Received:** January 03, 2024; **Accepted:** January 17, 2024; **Published:** January 23, 2024

**Copyright:** © 2024 Simitopoulou M, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

not precisely targeted to the particular health issues historically encountered on a specific farm or group of farms. Instead, they tend to be added to the feed or water as a matter of routine by feed manufacturers or farmers with very little prior discussion with the farm veterinarian on the best choice for the particular farm circumstances. In this study ten commercial broiler farms, in Cyprus and Greece, with relatively high historic reliance on antibiotic use were selected to be monitored for both antibiotic usage and biological performance, before (pre-additive production cycles) and after (post-additive production cycles) the targeted application of feed and/or water NAAs. Prior to this, the same farms were also involved in biosecurity improvements implemented after audits carried out following the Wagenigen BiosEcurity Assessment Tool (BEAT). The post-additive production cycles compared to the pre-additive production cycles indicated that NAA use was accompanied by a 50.3% decrease in the number of antibiotic treatments ( $p < 0.01$ ), a non-significant 21.4% reduction in the days of treatment as well as a 5.7% improvement in biological performance as measured by the European Production Efficiency Factor (EPEF) ( $p < 0.1$ ). More substantial improvements were also noted when the post-additive production cycles were compared with the pre-additive production cycles before the biosecurity improvements implementation. In this case the antibiotic treatments were shown to have decreased by 58.7% ( $p < 0.01$ ), and this was accompanied by a 52% decrease in the days of treatment ( $p < 0.01$ ) as well as an improvement of 16.9% ( $p < 0.01$ ) in biological performance as measured by EPEF. On the basis of this work, it may be concluded that the targeted use of feed and/or water additives when based on the specific farm health issues historically encountered can improve the biological performance and in parallel reduce the use of antibiotics in broiler farms.

**Keywords:** Antibiotics; Broilers; Commercial farms; Feed additives; Performance

## Introduction

The use of antibiotics as growth promoters (AGPs) in animal feeds has been permitted in the member states of the European Union (EU) during decades. However, concerns about the development of antimicrobial resistance (AMR) and the spreading and transfer of antibiotic resistance genes from animal to human sectors, led to the withdrawal of the approval for AGPs in the EU since January 1st, 2006 [1]. Available data suggest that although resulting in a reduction of the overall use of antibiotics in animals, the AGP ban in Europe has associated with a substantial increase in the use of therapeutic antibiotics for food animals [2]. In the poultry sector, there is evidence that the initial removal of AGPs has been associated in some cases to poultry performance problems, feed conversion increases and the rise of certain poultry diseases particularly enteric such as (subclinical) necrotic enteritis [3]. The increasing sociopolitical concerns regarding the development of AMR as well as at the same time the need to prevent farmer's economic losses has led to the development of alternatives to antibiotics. The primary goal of these alternative methods is disease prevention and reduction in veterinary intervention in commercial livestock farming, in order to improve animal growth performance, while based on the safety, efficacy and cost-effectiveness of such

Natural Antibiotic Alternatives (NAAs) [4,5]. The term NAA includes probiotics, prebiotics, symbiotics, organic acids, essential oils, enzymes, immunostimulants, and phytochemicals (phytobiotics) such as botanicals, herbs, essential oils, and oleoresins. These are the most common feed and/or water additives that are currently widely used in the poultry industry following the ban of AGPs. They can be easily mixed with other feed ingredients, have no tissue residues of concern, and are often shown to improve feed intake, weight gain, feed conversion rate and strengthen bird immunity as well as improve digestive processes. Overall, these effects have a potential positive action on animals' health and hence have the potential to decrease reliance on antibiotics [6]. Among the commercially available NAAs, the organic acids play an important role in gut health in animals. These acidifiers could be used to favorably impact the intestinal microbial populations and improve the immune response, hence perform an activity similar to antibiotics in food animals in countering pathogenic bacteria [7]. Short-chain organic acids have a specific antimicrobial activity that is pH dependent. Reductions in bacteria are associated with feeding organic acids, which are particularly effective against acid-intolerant species such as *E. coli*, *Salmonella*, and *Campylobacter*. Both antibiotics and organic acids improve protein and energy digestibility by reducing microbial competition with the host for nutrients and endogenous nitrogen losses, by lowering the incidence of subclinical infections and secretion of immune mediators, and by reducing production of ammonia and other growth-depressing microbial metabolites. Organic acids have several additional effects that go beyond those of antibiotics. These effects include reduction in digesta pH, increased pancreatic secretion, and trophic effects on the gastrointestinal mucosa [8].

Another NAA group consists of plant-based mixtures. These are also referred to as phytochemicals, phytobiotics or phytochemicals and are natural bioactive compounds that are derived from plants. A wide variety of herbs and spices (e.g., thyme, oregano, rosemary, marjoram, yarrow, garlic, ginger, green tea, black cumin, coriander and cinnamon) have been used in poultry for their potential application as AGP alternatives. In addition to herbs and spices, various essential oils and phytochemicals (thymol, carvacrol, cinnamaldehyde, eugenol, and extracts from coriander, star anise, ginger, garlic, rosemary, turmeric, basil, caraway, lemon and sage) have been used individually or as blends to improve animal health and performance. The beneficial effects of phytochemicals are attributed to their antimicrobial and antioxidant properties. In addition, the inclusion of phytochemicals in the diets alters and stabilizes the intestinal microbiota and reduces microbial toxic metabolites [9]. Exogenous enzymes such as xylanases and glucanases have been shown to improve feed utilization of dietary components such as protein, amino acids, starch, lipids, and energy. Feed enzymes can affect the gastrointestinal tract's microbial ecosystem by reducing undigested substrates and anti-nutritive factors and producing oligosaccharides *in situ* from dietary non-starch polysaccharides with potential prebiotic effects [10]. Probiotics are microorganisms which favor the healthy function of the animal's intestinal tract. A healthy intestinal tract has positive effects on the circulatory and immune system. Many different nutrients, such as pectin, cellulose and xylans, have been shown to promote the development of various favorable intestinal microorganisms and as such have a prebiotic effect. Prebiotics are not extensively metabolized, and have been shown to encourage targeted metabolic processes, which in turn bring health benefits to the animal's ecosystem [11]. It is common commercial practice that

feed manufacturers use a holistic approach in the choice of NAAs included in their poultry feed range. Such an approach is therefore not usually targeted on individual farm situations and nor does it take into account historic farm problems and veterinary interventions. It is also associated with few dialogues between the farmer, the feed supplier, or the farm veterinarian, about the choice of the feed and/or water additive best suited to the individual farm situations. The purpose of this field study was to test in commercial broiler farms from two countries, Cyprus and Greece, a set of NAAs which were selected after due consideration and discussion between the farm veterinarian establishing which were best suited to counter the specific problems identified from historical fact-based information that considered biological performance data, health data and veterinary diagnoses and consequent interventions.

## Materials and Methods

### Characteristics of the farms


Ten commercial broiler farms in two EU countries (Greece and Cyprus) were involved in this study. For the purposes of this study the farms were allocated to five groups AG, BG, AC, BC and CC depending on the historic information provided. All the farms were selected on the basis that they had a consistent and relatively high reliance on antimicrobial usage in their most recent growing cycles. As such, they were not selected to be representative of each country in either biological performance or antimicrobial usage. In Greece, five separate broiler farms with bird populations above 15,000 each, were enrolled. In Cyprus, five broiler farms each also with above 15,000 birds participated. All the farms had taken part shortly before to a separate study investigating the effect of health plan improvement implementations, post biosecurity audits carried out using BEAT (Base Editing Analysis Tool). BEAT is a risk analysis methodology developed by the University of Wageningen on the effects on biological performance and antimicrobial usage [12]. It is based on the FAO 3-Zone Biosecurity model, the BIOCheck UGENT [13] and the Dutch Hygiene scan. It has been described in full by Schreuder et al. [12].

The broiler houses were typical of intensive livestock facilities in the two countries, all with automatic feeders and drinkers and all tunnel ventilated. The heating systems were mostly air to air heat exchangers. The light intensity and photoperiodism followed a similar pattern in all cases. Straw or rice hull litter was used. Plant based diets were used throughout. These followed the nutrient recommendations as well as the feeding program as described in the Aviagen Ross 308 guidelines for birds targeted to be slaughtered at live weight 2.5 to 3.0 kg. The feed used was in the form of mash and was farm mixed from the grower stage onward while for the starter (0-10 days of age) commercially available crumbled broiler feed was used in both countries. Feed and water were offered *ad libitum* throughout the whole cycle.

The day-old chicks originated from one hatchery in each country throughout all the recorded periods. In all cases they were vaccinated at hatch, for Marek, Infectious Bronchitis and Newcastle Disease. No detailed information on the parent stock health statuses were made available. All animal care procedures complied with European Directives 2007/43/EC, 2005/1/EC and 2009/1099/EC which set welfare standards for keeping chickens for meat production, for their transport to the slaughterhouse and their protection during slaughter procedure.

## Time line of Study

The 4 periods described in (Figure 1) are the results of the successive implementations of two distinct procedures, namely the implementation of biosecurity measures (called Int or BioInt) and the implementation of Natural Antibiotic Alternatives (called Add). The 4 periods are as follows:



- The historic pre-intervention and pre-additive period (PreInt-PreAdd) corresponding to five consecutive production cycles, where the data collected and ascribed to those cycles was made available to the research team by the farmer and his veterinarian.
- The biosecurity improvement intervention period (BioInt), posterior to the biosecurity risk analysis (BEAT), covering two production cycles, where improvements were carried out.
- The pre-additive monitored period (PreAdd) also covering two production cycles, where details of biological performance as well as veterinary interventions were recorded by the research team. This period was then treated as the control.
- The similarly monitored post-additive production period (PostAdd) of the same duration which was considered as the treatment.

The biosecurity improvements were continuously implemented during the 3 last periods.

## Biological performances, health status of the farms and NAAs description and usage by farm

The information based on data recorded during the five production cycles corresponding to the PreInt-PreAdd periods was made available from all the participant farms. Such information included biological performance data covering average slaughter weight, average daily weight gain, feed conversion ratio (FCR), total mortality rate, footpad score (scored from 0 to 2), according to the scoring described by Berg [14] and the European Production Efficiency Factor (EPEF), where EPEF was calculated using the following formula:

$$EPEF = \frac{[[\text{Livability (\%)} \times \text{Live Weight (kg)}] / [\text{Age (days)} \times \text{FCR}]] \times 100$$

Antimicrobial usage records for the PreInt-PreAdd period were also made available, as was the veterinary diagnosis attached to each production cycle. These information bases provided the platform upon which discussions involving the farmer, his veterinary consultant and researchers participating in this study took place so as to arrive at an agreed individual farm-based additive regime for the

PostAdd period. All such individual farm-based additive regimes made use of commercially available and approved feed and/or water products, at best targeted at a possible reduction of antimicrobial use at each specific site, while also potentially able to maintain similar biological performance as measured by the parameters described above. The seven commercially available feed and/or water additives used in the study are described below in (Table 1).

Feed/Water Additive	Description
B-act® (Huvepharma®)	Probiotic feed additive containing viable spores of a unique strain of <i>Bacillus licheniformis</i> producing bacteriocins
Avipremium® D (Vetagro®)	Concentrated source of tributyrin in powder form (55% butyric acid).
Biotronic® Top 3 (DSM/ex Biomin product)	Combination of synergistically acting organic acids and flavoring compounds: Formic Acid 20% Acetic Acid 10% Propionic Acid 5%
Digestarom® P.E.P Sol (DSM/ex Biomin product)	Liquid phytogetic blend of natural extracts of aromatic plants containing: Mixture of flavoring compounds E484 Glyceryl polyethyleneglycol ricinoleate Propyleneglycol
Biotronic® Top Liquid (DSM/ex Biomin product)	Combination of synergistically acting organic acids and flavoring compounds

**Table 1: Feed and water additives used in this study.**

The (Table 2) depicts the feed and/or water additives used by farm and/or group of farms, the dosage and the administration route as well as the period of addition. As already mentioned, the 5 groups of farm(s)/additive(s) combinations were the results of the analysis of the PreInt-PreAdd information on both biological performance and veterinary interventions, and more importantly the disease challenges faced during this period. The AG and BG farms faced specific problems identified as omphalitis, necrotic enteritis and colibacillosis. The farms grouped as AC suffered from cases of dysbacteriosis, necrotic enteritis, peritonitis, arthritis and air sacculitis. Group BC suffered from omphalitis, necrotic enteritis, air sacculitis and arthritis while farm group CC suffered from necrotic enteritis, air sacculitis, peritonitis and arthritis. Groups AG and BG where the water source was farm wells also suffered from poor water quality.

	Number of	Feed / Water	Dosage	Period	Administration
	Broiler	Additive	(per 1000 kg of	(days)	Route
	Houses		feed or 1000 L		
			of water)		
Group AG	two of equal capacity	PoultryStar® ME	500	11 to 40	feed
		Biotronic® Top 3	1000	11 to 40	feed
		Biotronic® Top Liquid	1200	4 to 11	water

		Digestarom® P.E.P Sol	100	15 to 27	water
Group BG	three of equal capacity	AdiCox® AP	60	11 to 40	feed
		B-act®	500	11 to 40	feed
		Biotronic® Top Liquid	1200	4 to 11 and 16 to 28	water
Group AC	two of equal capacity	B-act®	500	10 to slaughter	feed
Group BC	two of equal capacity	Biotronic® Top 3	1000	10 to slaughter	feed
Group CC	one	Avipremium® D	200	10 to slaughter	feed

**Table 2:** Description of the NAAs used per Group.

## Statistical analysis

All the analyses were performed in R 4.1.0 (R Core Team, 2021). To assess the biological performance and antibiotic use during the PreAdd and PostAdd periods, mean scores were determined. To compare the results between the Pre-Add and Post-Add periods, and Analysis of Variance (ANOVA) test was performed. To assess the biological performance and antibiotic use during the PreInt-PreAdd and PostAdd periods, mean scores were also determined. To compare the results between the PreInt-PreAdd and PostAdd periods, Wilcoxon-Rank-Sum tests were performed. The differences were considered statistically significant with  $p < 0.05$ .

## Results

### Pre-Add and Post-Add comparison

The key biological performance indices as well as usage of antibiotic treatments are summarised in (Table 3). Overall, comparison of the PostAdd and the PreAdd periods showed a significant ( $p < 0.05$ ) reduction of the mortality rate (-37%), accompanied by a slight but not statistically significant improvement of EPEF (+5.7%;  $p < 0.1$ ). In addition, a strong reduction (-50%) of the average number of antibiotic treatments per production cycle was observed ( $p < 0.01$ ). It was accompanied by a non-significant parallel reduction in the overall days of antibiotic treatment (-21.4%;  $p > 0.2$ ).

### PreInt-PreAdd and PostAdd comparison

As the PreAdd period shown in (Table 3) was monitored subsequent to the implementation of Biosecurity Improvements (BioInt period) in the farms, this provided the opportunity to perform an additional comparison between the PreInt-PreAdd period - that is before biosecurity improvements and NAA addition -, and the Post-Add period when both biosecurity improvements and NAA addition are present. This comparison is summarised in (Table 4), showing that all indicators were significantly improved. In particular, EPEF value showed a highly significant improvement in the PostAdd period (+16.9%;  $p < 0.001$ ), emanating from improvements in feed conversion

	Mortality Rate	EPEF	Number of AB Treatments	Days of AB Treatments
PreAdd	4.22 ± 0.42	333 ± 7	1.91 ± 0.28	4.39 ± 0.58
PostAdd	2.65 ± 0.44	352 ± 8	0.95 ± 0.15	3.45 ± 0.57
p-value*	0.014	0.085	0.006	0.259
	$p < 0.05$	$p < 0.1$	$p < 0.01$	N.S.

**Table 3:** Average Biological Performance and Antibiotic Use for the Pre-Add and PostAdd Periods.

\*ANOVA Test

ratio, livability and final live weight (not shown). Similarly, both the number of antibiotic treatments as well as the days of antibiotic treatments per production cycle showed substantial and highly significant reductions: -58.7% ( $p < 0.001$ ) and -52% ( $p < 0.001$ ), respectively.

	Mortality Rate	EPEF	Number of AB Treatments	Days of AB Treatments
PreInt-PreAdd	4.04 ± 0.32	301 ± 7	2.30 ± 0.12	7.19 ± 0.52
PostAdd	2.65 ± 0.44	352 ± 8	0.95 ± 0.15	3.45 ± 0.57
p-value*	0.004882	0.000043	0.0000057	0.00012
	$p < 0.01$	$p < 0.001$	$p < 0.001$	$p < 0.001$

**Table 4:** Average Biological Performance and Antibiotic Use during the PreInt-PreAdd and Post Add periods.

\* Wilcoxon Test

## Discussion

The implementation of farm-tailored Non-Antibiotic Additives (NAAs) was tested in the present study in a context of overall improvement of healthy status of poultry farms. The implementation of farm-tailored Non-Antibiotic Additives (NAAs) was tested in the present study in a context of overall improvement of healthy status of poultry farms. In this context it is recognized that the most beneficial strategy for reducing the need for the use of antibiotic treatment under commercial field conditions is to prevent pathogens from entering and spreading in the farms [13-15]. This can best be done in practice by improving farm biosecurity culminating in the development and implementation of effective flock health and welfare plans. Schreuder et al. [12] discussed in detail how a farm can be divided into three zones which include the area outside the farm (red zone), the area and buildings within the farm except of the poultry houses (orange zone) and finally the poultry houses themselves (green zone). Each zone has its own parameters that need to be assessed on the basis of correct biosecurity. If those parameters don't meet with the proper standards, then they should be changed in order to improve the farms biosecurity. For example, the red zone should be checked for proper pest control as well as personnel, visitor and vehicle access protocols. As far as the orange zone is concerned, cadaver, manure, bedding and feed storages must be tightly sealed and isolated, and cleaning and disinfection of the farm yard as well as the buildings in the farm should be done regularly. Finally, in the green zone cleaning and disinfection of both the poultry house and the entry room and the immediate removal of dead birds are of utmost importance. All the farms participating in this study went through this process before any NAA addition was targeted and used. The NAAs used in each group were chosen according to the groups'

specific problems identified from historical fact-based information that covered biological performance data, health data and veterinary diagnoses. For example, certain broiler houses had issues with dysbacteriosis, necrotic enteritis and arthritis. In this case B-act® was the feed additive of choice. Certain probiotics have the ability to suppress the effects of dysbacteriosis in broilers and help inhibit the growth of *Clostridium perfringens*, the direct causative agent of necrotic enteritis. This leads to a reduction in the flock's mortality, an improved average weight and a decrease in the feed conversion ratio which are findings that coincide with the results of this study. Another benefit of the use of probiotics is the reduction in the incidence of arthritis-induced lameness which is brought about by bacterial translocation from the intestines to the joints [16]. A common characteristic of other grouped broiler houses was the poor quality of water coupled with incidences of necrotic enteritis and arthritis. The feed additive chosen in this case was Biotronic® Top 3. Multiple previous studies have shown that inclusion of organic acids in the feed can lead to a significantly higher final weight, weight gain, and better feed/gain ratio [17, 18]. Organic acids can reduce the number of pathogenic bacteria and their entry into the intestinal mucosa and subsequently the inflammatory processes that will ultimately affect performance [19]. Moreover, the use of organic acids has been shown to significantly increase the height of villi from all segments of the small intestine and to cause a moderate to marked increase of the goblet cell counts, compared to not supplemented controls. Goblet cells are distributed along the villi and the mucin they produce plays a key role in the proper function of the intestinal epithelium [20]. In another group of broiler houses recording similar problems, the targeted NAA regime was based on the addition of Avipremium® D which contained the organic triglyceride tributyrin. There is increasing evidence suggesting that butyrate can act as the major energy source of the colon, reduce pathogen colonization, improve intestinal barrier function, exert anti-inflammatory properties and protect the birds against liver and kidney dysfunction. Dietary supplementation with tributyrin has been demonstrated to improve animal performance by increasing the final weight and decreasing the FCR in the entire growth phase. This improvement in performance is brought about by the proliferation and differentiation of intestinal mucosal cells which enhances nutrient absorption and utilization [21]. A different study has shown that dietary sodium butyrate supplementation can improve the growth performance in chickens under stress and that this may be used to moderate the immune response and reduce tissue damage with diseases such as necrotic enteritis, arthritis and peritonitis [22], as it was also the case of certain groups of farms in this study. It is often the case that some commercial farms and/or individual broiler houses within a farm encounter a number of health issues arising out of a complexity of challenges which extend from poor water quality or shortcomings in biosecurity, the latter being not easy to correct in the short term. In this study the targeted choice of additives was extended to both water and/or feed products with a broad range of claimed benefits. Specifically, one group of broiler houses received a combination of PoultryStar® ME which is a symbiotic product blend, together with the organic acid Biotronic® Top 3 in the feed, and Biotronic® Top Liquid and Digestarom® P.E.P Sol in the drinking water. The use of organic acids in both the feed and the water regimes was based on the premise that this combination would provide an enhanced control of Gram-negative pathogens, as well as a direct support to the healthy gut microbiota in the face of a more severe pathogen challenge. Davidson et al. [23] reported that when the bacterial internal pH drops because of the presence of an organic acid, the pathogenic bacteria

must use adenosine triphosphate to actively transport excess protons from the interior of the cell, resulting in a depletion of cellular energy and cell death. Another possible reason for the modification of intestinal microbial growth by organic acids can pertain to their prebiotic effects, that result in the production of short-chain fatty acids from bacteria which are also thought to have an antibacterial effect by decreasing luminal pH [24].

The targeted use of the symbiotic mix in this study was also based on the recorded observation that the birds in this group of farms showed frequently increased incidences of lameness attributed to arthritis as well as poor litter quality. According to the study of Stark [25], by combining a pro- with a prebiotic a significant overall reduction in the onset of lameness attributed to bacterial chondronecrosis was observed in comparison with control broilers receiving the non-supplemented diet alone (19.6% vs. 33.2%, respectively). This effect was attributed to the probiotic eliciting beneficial responses to the bird's gastrointestinal microflora and immune system, contributing in reducing or stopping bacterial translocation from the gastrointestinal tract to susceptible leg joints. Furthermore, another study [26] indicated that these types of symbiotic products were able to prevent clinical signs associated with *Eimeria* challenge in poultry, and alleviate performance losses when included in the diet, especially regarding oocyst shedding. The increase in EPEF as well as the decrease in the FCR observed in the present study agrees with a study carried out by Awad et al. [27] who found that the dietary supplementations of pre- and probiotics resulted in an increase in the villus height and crypt depth ratio of intestinal mucosa of broilers. Subclinical coccidiosis, particularly where poor litter quality is encountered, is often a problem leading to additional enteric stresses such as dysbacteriosis and necrotic enteritis. This is particularly a problem where the system does not follow 'an all-in all-out' practice, and bed collection at depletion is carried out over a period of time. Blends of herbs and spices as well as essential oils and other plant extracts have been used widely in the industry to help coccidiosis control. In this study where such practices and historic problems were encountered, the targeted NAA approach used was based on such products namely Digestarom® P.E.P Sol and AdiCox® AP. Both products are plant-based preparations with reported coccidiostat efficacy. Murugesan et al. [28] reported that phytogenic feed additives support the establishment of a favorable gut microbiota composed of higher numbers of *Lactobacillus* spp. and fewer *Clostridium* spp. Furthermore, the same study found that supplementation of an herb-based mixture to a coccidiostat free diet increased the body weight gain and lowered the FCR.

AdiCox® AP has specifically been shown to act as a natural coccidiostat [29]. This coincides with the results of the present study in which mortality was decreased and the EPEF was increased significantly when using these types of phytogenic products on farms with a history of coccidiosis challenges. The present findings confirm that commercially available NAAs when supported by both an explanation of the mode of action and published efficacy results, can provide a very useful approach in an effort to further reduce the need of veterinary interventions in the form of metaphylaxis and certainly prophylaxis in broiler production. As individual feed and/or water additives often have a different mode of action and application target, the efficacy of such use lies to a large degree on good knowledge of the historic problems and health issues prevailing on the targeted farm. The present study has clearly demonstrated this in the reduction of antibiotic usage after the targeted NAAs addition.

Biosecurity is a major issue facing many commercial broiler farms, particularly those situated in heavily populated poultry areas. Improvements in biosecurity are possible through training and personnel involvement as well as through capital investment, resulting in biological improvements and less reliance on veterinary interventions based on antibiotic use [12]. The present study has shown that when such biosecurity improvements are associated with targeted NAA solutions, even greater improvements in both antibiotic reduction and biological performances can be demonstrated.

## Conclusion

Modern poultry production systems often constrain natural poultry behavior and may result in health problems which in turn necessitate the use of antibiotics under veterinary prescription. Within the European Union there is increased pressure to limit if not prohibit the use of antibiotics for prophylactic as well as metaphylactic purposes in food producing animals. Decreased reliance on antibiotics can be achieved by improving farm biosecurity. For this to be effective a detailed and planned biosecurity audit involving both the farmer and the veterinarian across a structured protocol is a prerequisite. Further reduction in antibiotic reliance can be achieved through the targeted selection and use of a non-antibiotic feed and/or water commercially available products best targeted on the specific farm historic health issues.

## Acknowledgement

This research was part of the EU-China Healthy Livestock project. The authors wish to acknowledge that Healthy Livestock is funded by the European Union H2020 research and innovation program under grant agreement number 773436. The European Commission's support for the production of this publication does not constitute an endorsement of the contents, which reflect the views only of the authors, and the Commission cannot be held responsible for any use which may be made of the information contained therein. BEAT tool project was co-funded by the Dutch Ministry of agriculture, nature and food quality via the TKI Agri & Food under grant agreement number AF-EU18020.

## Conflicts of Interest

The authors declare no conflicts of interest.

## References

1. Castanon JIR (2007) History of the Use of Antibiotic as Growth Promoters in European Poultry Feeds. *Poult Sci* 86: 2466-2471.
2. Casewell M, Marco E, Friis C, McMullin P, Phillips I (2003) The European ban on growth-promoting antibiotics and emerging consequences for human and animal health. *J Antimicrob Chemother* 52: 159-161.
3. Dibner JJ, Richards JD (2005) Antibiotic growth promoters in agriculture: history and mode of action. *Poult Sci* 84: 634-643.
4. Shivaramaiah S, Pumford NR, Morgan MJ, Wolfenden RE, Wolfenden AD, et al. (2011) Evaluation of *Bacillus* species as potential candidates for direct-fed microbials in commercial poultry. *Poult Sci* 90: 1574-1580.
5. Papatsiros VG, Katsoulou PD, Koutoulis KC, Karatzia M, Dedousi A et al. (2013) Alternatives to antibiotics for farm animals. *CABI Rev* 1-15.
6. Abd El-Hack ME, El-Saadony MT, Salem HM, El-Tahan AM, Soliman MM, et al. (2022) Alternatives to antibiotics for organic poultry production: types, modes of action and impacts on bird's health and production. *Poult Sci* 101: 101696.
7. Pearlín BV, Muthuvel S, Govidasamy P, Villavan M, Alagawany M, et al. (2020) Role of acidifiers in livestock nutrition and health: A review. *J Anim Physiol Anim Nutr* 104: 558-569.
8. Dibner JJ, Buttin P (2002) Use of Organic Acids as a Model to Study the Impact of Gut Microflora on Nutrition and Metabolism. *J Appl Poult Res* 11: 453-463.
9. Lillehoj H, Liu Y, Calsamiglia S, Fernandez-Miyakawa ME, Chi F, et al. (2018) Phytochemicals as antibiotic alternatives to promote growth and 401 enhance host health. *Vet Res* 49: 76.
10. Anadón A, Ares I, Martínez-Larrañaga MR, Martínez MA (2019) Enzymes in Feed and Animal Health, in: Gupta, R.C., Srivastava, A., Lall, R. (Eds.), *Nutraceuticals in Veterinary Medicine*. Springer Int Publ Cham 303-313.
11. Markowiak P, Śliżewska K (2018) The role of probiotics, prebiotics and synbiotics in 407 animal nutrition. *Gut Pathog* 10: 21.
12. Schreuder J, Simitopoulou M, Angastiniotis K, Ferrari P, Wolthuis-Fillerup M, et al. (2023) Development and implementation of a risk assessment tool for broiler farm biosecurity and a health intervention plan in the Netherlands, Greece, and Cyprus. *Poult Sci* 102: 102394.
13. Gelaude P, Schlepers M, Verlinden M, Laanen M, Dewulf J (2014) Biocheck. UGent: A quantitative tool to measure biosecurity at broiler farms and the relationship with technical performances and antimicrobial use. *Poult Sci* 93: 2740-2751.
14. Berg C (1998) Foot-pad dermatitis in broilers and turkeys. Dr. Thesis.
15. Newell DG, Elvers KT, Dopfer D, Hansson I, Jones P, et al. (2011) Biosecurity- Based Interventions and Strategies To Reduce *Campylobacter* spp. on Poultry Farms. *Appl Environ Microbiol* 77:8605-8614.
16. Xu T, Yue K, Zhang C, Tong X, Lin L, et al. (2022) Probiotics Treatment of Leg Diseases in Broiler Chickens: a Review. *Probiotics and Antimicro Prot* 14: 415-425.
17. Agbai KN, Omega JJ, Sekoni AA, Afolayan M (2020) Effect of Different Levels of Biotronic® Top Forte (Commercial blended organic acids) as a Replacement for Antibiotic Growth Promoter on the Growth Performance and Villi Morphology of Pullet Chicks (0-8weeks). *Niger J Anim Sci Technol NJAST* 3: 13-21.
18. Sara A, Marti R, Bentea MI, Toader I, Gabor EF (2016) The Effects of Some Prebiotic Products (Actigen, Biotronic Top3) Administrated in The Food of Broiler Chickens on the Meat Quality. *Bull Univ Agric Sci Vet Med Cluj-Napoca Anim Sci Biotechnol* 73: 202-433.
19. Khan RU, Naz S, Raziq F, Qudratullah Q, Khan NA, et al. (2022) Prospects of organic acids as safe alternative to antibiotics in broiler chickens diet. *Environ Sci Pollut Res* 29: 32594-32604.
20. Kum S, Eren U, Onol AG, Sandikci M (2010) Effects of dietary organic acid supplementation on the intestinal mucosa in broilers. *Rev Med Vet* 161: 463-468.
21. Gong L, Xiao G, Zheng L, Yan X, Qi Q, et al. (2021) Effects of Dietary Tributyrin on Growth Performance, Biochemical Indices, and Intestinal Microbiota of Yellow-Feathered Broilers. *Anim. Open Access J. MDPI* 11: 3425.
22. Zhang WH, Jiang Y, Zhu QF, Gao F, Dai SF (2011) Sodium butyrate maintains growth performance by regulating the immune response in broiler chickens. *Br Poult Sci* 52: 292-301.
23. Davidson PM, Taylor TM, Schmidt SE (2014) Chemical Preservatives and Natural Antimicrobial Compounds, in: Doyle, MP, Buchanan RL (Eds.), *Food Microbiology*. ASM Press 765-801.
24. Corrier DE, Hinton A, Ziprin RL, Beier RC, DeLoach JR (1990) Effect of Dietary Lactose on Cecal pH, Bacteriostatic Volatile Fatty Acids, and *Salmonella typhimurium* Colonization of Broiler Chicks. *Avian Dis* 34: 617.
25. Stark JM (2013) Efficacy of Probiotics for Reducing the Incidence of Lameness in Broilers Grown on Wire Flooring. Graduate Theses and Dissertations.

26. Duff AF, Briggs WN, Chasser KM, Lilburn MS, Syed B, (2020) Effect of dietary synbiotic supplementation on performance parameters in turkey poults administered a mixed *Eimeria* species inoculation I *Poult Sci* 99: 4235-4241.
27. Awad WA, Ghareeb K, Abdel-Raheem S, Böhm J (2009) Effects of dietary inclusion of probiotic and synbiotic on growth performance, organ weights, and intestinal histomorphology of broiler chickens. *Poult Sci* 88: 49-56.
28. Murugesan GR, Syed B, Haldar S, Pender C (2015) Phytogetic Feed Additives as an Alternative to Antibiotic Growth Promoters in Broiler Chickens. *Front Vet Sci* 2.
29. Łukasiewicz M, Pietrzak D, Michalczuk MJ (2013) Effect of selected feed additives used in coccidiosis prophylaxis on production results and meat quality of fast-growing Hubbard Flex chickens. *Arch. Geflügelkunde*.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>